

# **AMCS / CS 247 – Scientific Visualization**

## **Lecture 29: Vector Field / Flow Visualization, Pt. 8**

Markus Hadwiger, KAUST

# Reading Assignment #15++




Read (required):

- Go over the flow visualization reading assignments again

# Lagrangian vs. Eulerian



- Lagrangian: move along with the particle
- Eulerian: consider fixed point in space, look at particles moving through



- Example for pixels: rotate image (a),  
Lagrangian: move pixels forward (b),  
Eulerian: fetch pixels from backward direction (c)

# Material Derivative (1)



The material derivative (convective derivative) gives the rate of change when following a particle in the flow

$$\frac{D}{Dt} = \frac{\partial}{\partial t} + (\mathbf{u} \cdot \nabla)$$

# Material Derivative (1)



The material derivative (convective derivative) gives the rate of change when following a particle in the flow

$$\frac{D}{Dt} = \frac{\partial}{\partial t} + (\mathbf{u} \cdot \nabla)$$

$$\frac{DT}{Dt} = \frac{\partial T}{\partial t} + (\mathbf{u} \cdot \nabla) T$$

# Material Derivative (1)



The material derivative (convective derivative) gives the rate of change when following a particle in the flow

$$\frac{D}{Dt} = \frac{\partial}{\partial t} + (\mathbf{u} \cdot \nabla)$$

$$\frac{DT}{Dt} = \frac{\partial T}{\partial t} + (\mathbf{u} \cdot \nabla) T$$

$$\frac{DT}{Dt} = \frac{\partial T}{\partial t} + u \frac{\partial T}{\partial x} + v \frac{\partial T}{\partial y} + w \frac{\partial T}{\partial z}$$

## Material Derivative (2)



Actually, nothing else than application of the multi-variable chain rule:

$$dT = \frac{\partial T}{\partial t} dt + \frac{\partial T}{\partial x} dx + \frac{\partial T}{\partial y} dy + \frac{\partial T}{\partial z} dz$$

# Material Derivative (2)



Actually, nothing else than application of the multi-variable chain rule:

$$dT = \frac{\partial T}{\partial t} dt + \frac{\partial T}{\partial x} dx + \frac{\partial T}{\partial y} dy + \frac{\partial T}{\partial z} dz$$

$$dT = \frac{\partial T}{\partial t} dt + \frac{\partial T}{\partial x} \frac{dx}{dt} dt + \frac{\partial T}{\partial y} \frac{dy}{dt} dt + \frac{\partial T}{\partial z} \frac{dz}{dt} dt$$



# Material Derivative (2)



Actually, nothing else than application of the multi-variable chain rule:

$$dT = \frac{\partial T}{\partial t} dt + \frac{\partial T}{\partial x} dx + \frac{\partial T}{\partial y} dy + \frac{\partial T}{\partial z} dz$$

$$dT = \frac{\partial T}{\partial t} dt + \frac{\partial T}{\partial x} \frac{dx}{dt} dt + \frac{\partial T}{\partial y} \frac{dy}{dt} dt + \frac{\partial T}{\partial z} \frac{dz}{dt} dt$$

$$\frac{dT}{dt} = \frac{\partial T}{\partial t} + \frac{\partial T}{\partial x} \frac{dx}{dt} + \frac{\partial T}{\partial y} \frac{dy}{dt} + \frac{\partial T}{\partial z} \frac{dz}{dt}$$

# Material Derivative (2)



Actually, nothing else than application of the multi-variable chain rule:

$$dT = \frac{\partial T}{\partial t} dt + \frac{\partial T}{\partial x} dx + \frac{\partial T}{\partial y} dy + \frac{\partial T}{\partial z} dz$$

$$dT = \frac{\partial T}{\partial t} dt + \frac{\partial T}{\partial x} \frac{dx}{dt} dt + \frac{\partial T}{\partial y} \frac{dy}{dt} dt + \frac{\partial T}{\partial z} \frac{dz}{dt} dt$$

$$\frac{dT}{dt} = \frac{\partial T}{\partial t} + \frac{\partial T}{\partial x} \frac{dx}{dt} + \frac{\partial T}{\partial y} \frac{dy}{dt} + \frac{\partial T}{\partial z} \frac{dz}{dt}$$

$$u := \frac{dx}{dt}, \quad v := \frac{dy}{dt}, \quad w := \frac{dz}{dt}$$

# Material Derivative (2)



Actually, nothing else than application of the multi-variable chain rule:

We are given  $T(x, y, z, t)$  with four independent variables;

But now we want to go along a parameterized path with parameter  $t$ ,  
so  $x, y, z$  become dependent variables:  $x(t), y(t), z(t)$

Along this path, our goal is now to compute the derivative of the function

$T(x(t), y(t), z(t), t)$  with  $t$  as only independent variable:

$$\frac{d}{dt}T(x(t), y(t), z(t), t) =$$
$$\frac{\partial}{\partial t}T(x, y, z, t) + \frac{\partial}{\partial x}T(x, y, z, t) \frac{d}{dt}x(t) + \frac{\partial}{\partial y}T(x, y, z, t) \frac{d}{dt}y(t) + \frac{\partial}{\partial z}T(x, y, z, t) \frac{d}{dt}z(t)$$

$$u(t) := \frac{dx(t)}{dt}, \quad v(t) := \frac{dy(t)}{dt}, \quad w(t) := \frac{dz(t)}{dt}$$

# Advection



Advection equation; velocity field  $\mathbf{u}(x, y, z, t)$ ,  
no change following particle, just advection:  
set material derivative = 0:

$$\frac{\partial T}{\partial t} + (\mathbf{u} \cdot \nabla) T = 0$$

In the Navier-Stokes equations: “self-advection” of velocity

- Advect scalar components of velocity field individually  
(actually two equations in 2D, three equations in 3D)

$$\frac{\partial \mathbf{u}}{\partial t} = -(\mathbf{u} \cdot \nabla) \mathbf{u}$$

# Fluid Simulation: Navier Stokes (1)



Incompressible (divergence-free) Navier Stokes equations

$$\frac{\partial \mathbf{u}}{\partial t} = -(\mathbf{u} \cdot \nabla) \mathbf{u} - \frac{1}{\rho} \nabla p + \nu \nabla^2 \mathbf{u} + \mathbf{F},$$

$$\nabla \cdot \mathbf{u} = 0,$$

Components:

- Self-advection of velocity (i.e., advection of velocity according to velocity)
- Pressure gradient (force due to pressure differences)
- Diffusion of velocity due to viscosity (for viscous fluids, i.e., not inviscid)
- Application of (arbitrary) external forces, e.g., gravity, user input, etc.

## Fluid Simulation: Navier Stokes (2)



Actually, the momentum equation is a system of equations  
(2 equations in 2D, 3 equations in 3D)

For 2D:

these are PDEs!

$$\frac{\partial u}{\partial t} = -(\mathbf{u} \cdot \nabla) u - \frac{1}{\rho}(\nabla p)_x + \nu \nabla^2 u + f_x,$$

$$\frac{\partial v}{\partial t} = -(\mathbf{u} \cdot \nabla) v - \frac{1}{\rho}(\nabla p)_y + \nu \nabla^2 v + f_y.$$

# Thank you.

Thanks for material

- Ronny Peikert
- Helwig Hauser
- Meister Eduard Groeller
- Jens Krüger