

# **AMCS / CS 247 – Scientific Visualization**

## **Lecture 9: Scalar Fields, Pt. 3**

Markus Hadwiger, KAUST

# Reading Assignment #6 (until Mar 11)



Read (required):

- Real-Time Volume Graphics, Chapter 1 (*Theoretical Background and Basic Approaches*), from beginning to 1.4.4 (inclusive)
- Real-Time Volume Graphics, Chapter 2 (*GPU Programming*)

Optional: Refresh your memory on

- Partial derivatives of multivariate functions
- The concept of the gradient of a scalar-valued function
- The dot product between two vectors

# Quiz #1: Mar 4



## Organization

- First 30 min of lecture
- No material (book, notes, ...) allowed

## Content of questions

- Lectures (both actual lectures and slides)
- Reading assignments (except optional ones)
- Programming assignments (algorithms, methods)
- Solve short practical examples

# Gradient and Directional Derivative



Gradient  $\nabla f(x, y, z)$  of scalar function  $f(x, y, z)$ : (in Cartesian coordinates)

$$\nabla f(x, y, z) = \left( \frac{\partial f(x, y, z)}{\partial x}, \frac{\partial f(x, y, z)}{\partial y}, \frac{\partial f(x, y, z)}{\partial z} \right)^T$$

Directional derivative in direction  $\mathbf{u}$  :

$$D_{\mathbf{u}}f(x, y, z) = \nabla f(x, y, z) \cdot \mathbf{u}$$

And therefore also:

$$D_{\mathbf{u}}f(x, y, z) = \|\nabla f\| \|\mathbf{u}\| \cos \theta$$

# Gradient and Directional Derivative



Gradient  $\nabla f(x, y, z)$  of scalar function  $f(x, y, z)$ : (in Cartesian coordinates)

$$\nabla f(x, y, z) = \left( \frac{\partial f(x, y, z)}{\partial x}, \frac{\partial f(x, y, z)}{\partial y}, \frac{\partial f(x, y, z)}{\partial z} \right)^T$$

With basis vectors (Cartesian basis):

$$\nabla f(x, y, z) = \frac{\partial f(x, y, z)}{\partial x} \mathbf{i} + \frac{\partial f(x, y, z)}{\partial y} \mathbf{j} + \frac{\partial f(x, y, z)}{\partial z} \mathbf{k}$$

# The Dot Product (Scalar / Inner Product)



Angle between two vectors times their lengths


$$\mathbf{a} \cdot \mathbf{b} = \sum_{i=1}^n a_i b_i$$

$$\mathbf{a} \cdot \mathbf{b} = \|\mathbf{a}\| \|\mathbf{b}\| \cos \theta$$

(standard inner product in Cartesian coordinates)

Many uses:

- Project vector onto another vector,  
project into basis,  
project into tangent plane,  
...



# Thank you.

## Thanks for material

- Helwig Hauser
- Eduard Gröller
- Daniel Weiskopf
- Torsten Möller
- Ronny Peikert
- Philipp Muigg
- Christof Rezk-Salama